

# MUSTELID & VIVERRID CONSERVATION


The Newsletter of the IUCN/SSC  
Mustelid & Viverrid Specialist Group.

Number 1

August 1989



Liberian mongoose (*Liberiictis kuhni*). First photograph taken of a live animal. Photo by M. E. Taylor.



This number was produced with the support of the "Royal Zoological Society of Antwerp", Antwerp, Belgium, and with the kind assistance of Dr. G. M. Carpaneto, Dipartimento di Biologia animale e dell'uomo, Universita di Roma "La Sapienza", Roma, Italy.

# **Mustelid & Viverrid Conservation**

The Newsletter of the IUCN/SSC Mustelid & Viverrid Specialist Group.

**Editor-in-chief:** Harry Van Rompaey, Edegem, Belgium

**Editors:** Michael Riffel, Karlsdorf, FRG  
Arnd Schreiber, Heidelberg, FRG  
Roland Wirth, München, FRG

The views expressed in this Newsletter are those of the authors and do not necessarily reflect those of the IUCN, nor the IUCN/SSC Mustelid & Viverrid Specialist Group.

We are particularly grateful to Walter Rasmussen for reading the manuscripts and improving the English style.

The aim of this Newsletter is to offer the members of the IUCN/SSC M&VSG, and those who are concerned with mustelids or/and viverrids, brief papers, news items, abstracts, and titles of recent literature.  
All readers are invited to send material to:

---

**Mustelid & Viverrid Conservation**  
**c/o Dr. H. Van Rompaey**  
**Jan Verbertlei, 15**  
**2520 Edegem**  
**Belgium**

---

# EDITORIAL

This is the first number of what we hope will be a regular Newsletter series. That this Newsletter has surfaced is mainly due to the dedication and enthusiasm of Harry Van Rompaey who has offered to take over the major work in putting the publication together and have it sent out at regular intervals.

The aim of this Newsletter -like of all SSC Specialist Group Newsletters- is, of course, not to publish just another periodical for its own sake, but to intensify our efforts for the conservation of the species we are concerned with. As such the Newsletter wants to reach all who have some sort of interest in the survival of the present diversity of mustelids and viverrids. With animals scientifically so poorly known as many of the world's mustelids and viverrids we would like to see intensified efforts to learn more about the status, ecology, reproduction, diseases, and captive management needs of especially those species and subspecies listed as of conservation concern in the recently published Conservation Action Plan of the IUCN/SSC Mustelid & Viverrid Specialist Group.

As such the Newsletter wants to bring all persons together who want to work towards these aims in one way or another. We would appreciate to hear from anybody who would like to contact somebody to do conservation-related mustelid/viverrid research or who can offer his services and experience in starting one of the many priority actions identified in the Action Plan. We would be more than happy to publish such requests or offers in forthcoming issues of the Newsletter.

A major goal of the Newsletter will also be to give up- dates on recent conservation or research initiatives and to report on new or so far unrecognized conservation problems. If -as we hope-we will receive many short communications for this section, the Newsletter can quickly

fulfil its need to help coordinate action for threatened taxa and allow quick response to newly surfacing conservation problems.

While, as mentioned before, the Newsletter has become a reality thanks mainly to the efforts of Harry Van Rompaey, it needs the active cooperation of ALL MVSG members and it should not be too much of a request to ask at least one communication a year from each of you. Obviously anybody else who is not an official member of the Group and who has to say anything related to mustelid or viverrid conservation or the activities of the Specialist Group should also write.

Each member of the Specialist Group will receive a complementary copy of each Newsletter issue, but others interested in receiving it should pay something. In fact, the problem of funding any issues of the Newsletter additional to this present one is completely unsolved. It is your-the reader's-responsibility in helping to obtain a funding basis for the Newsletter. Initially we plan to produce two Newsletters a year, but to make the publication relatively attractive and to have enough surplus copies to be sent free of charge to researchers, students, government officials, etc. in third world countries we would need a minimum of US\$ 2,000 a year. I would suggest that any non-member of the Specialist Group who would like to subscribe to the Newsletter pays an annual fee of US\$ 15 (larger donations would, of course, be welcome). If each member of the Specialist Group and any other person interested in mustelid & viverrid conservation would find us four or five subscribers, the future of the Newsletter would be financially ensured.

**Roland Wirth**  
Chairman IUCN/SSC  
Mustelid & Viverrid Specialist Group

# Mustelidae and Viverridae from north-eastern Zaire: ethnozoological research and conservation.

Guiseppe M. CARPANETO and Francesco P. GERMI

The faunal richness of north-eastern Zaire is mainly due to the great variety of its vegetation types: lowland- and mountain forests, arid and wet savannahs, marsh and riverine ecosystems, etc. The high diversity of mammal communities is also found in the large number of small carnivore species. As regards the two families Mustelidae and Viverridae (including Herpestinae), they are respectively represented by 6 and 17 species.

In order to realize our own ethnozoological research on the forest- and savannah dwelling people (hunter-gatherers and shifting cultivators), we have carried out faunistic investigations on the mammals of the Kivu and Haut-Zaire regions. In the following checklist we marked (!) the species directly observed by us, at different times between 1984 and 1988; unmarked species were recorded by other authors in previous papers. Owing to taxonomists' disagreement through literature, we neglected the subspecies level. The following abbreviations were used: TM, Tshopo-Maiko Rivers region; IF, Ituri Forest; VI, Virunga National Park (including Mont Hoyo); GA, Garamba National Park.

Up to the present our ethnozoological surveys have been mainly carried out in the Ituri Forest and in the Virunga National Park.

## Ituri Forest

All the larger mammals serve as food for the Mbuti pygmies of the Ituri Forest. In this area the Alexander's cusimanse, *Crossarchus alexandri*, is the commonest carnivore species (Fig. 1), and is usually captured and killed by dogs but also often flushed out by dogs whereupon the Mbuti kill it with a stick blow on the head or shoot it with iron-tipped arrows. If the animal takes refuge in a hole, the Mbuti smoke it out.

During four consecutive bow-hunting days we observed a single Mbuti band catch four cusimanses. This species seems to be very common game for the Mbuti archers in the eastern Ituri Forest. On the other hand the Mbuti net-hunters of the western Ituri Forest do not have many chances of capturing cusimanses.

The above mentioned hunting techniques are used to kill the ratel (*Mellivora capensis*), the black-legged mongoose (*Bdeogale nigripes*), who are very aggressive against both dogs and men, *Herpestes* sp., and all the genet species.

The African civet (*Civettictis civetta*) and the two-spotted palm civet (*Nandinia binotata*) are often caught and eaten by the Mbuti. The former is easily captured with nets for antelopes because of its large size and terrestrial



Fig.1. Alexander's cusimanse (*Crossarchus alexandri*)

Photo G.M. Carpaneto



Fig.2. Mbuti girl wearing a hat made from *Genetta* skin.  
Photo G.M. Carpaneto

behaviour; the latter is killed with every kind of arrow (simple wooden, poisoned, and iron-tipped) when resting in trees.

Otters are widespread and common in the Ituri Forest but are only occasional game for pygmies, who lack hunting techniques for aquatic animals. Out of the water, when found near river banks, the otters may be roused by dogs and killed with iron-tipped arrows.

The skins of genets and civets are used to make hats. The Mbuti wear these hats when they go hunting for monkeys: they think that monkeys come near out of curiosity and so may be easily killed. The same hats are worn by Mbuti girls during their initiation ceremony (Fig. 2). The skin may also be used to make the wrist-protector bracelet which guards the archer's arm against the snap of the bow-string. The ratel's skin, on the other hand, is used to make drums.

Not only pygmies but also Bantu and Sudanic shifting cultivators inhabit the Ituri Forest. These people belong to

different tribes (Bira, Bali, Ndaka, Budu, Lese, etc.) and have close barter relations with Mbuti, who provide them with forest products such as meat and honey. These villagers, in fact, are not specialized hunters and view the forest as powerful and alien. They prefer to provide the Mbuti with starch foods and receive meat in exchange. Nevertheless they show an interesting trapping culture to integrate their protein requirements. The African civet and all mongooses are lured with bananas, oil palm fruits, and dead snakes as bait. The villagers appreciate eating all species of small carnivores (excluding their own clan totems) and use the animal skins in a similar fashion as the pygmies.

Several kinds of food restrictions exist in the traditional culture of both pygmies and villagers. For instance, according to the Mbuti of Alipanda (W. Ituri), the giant genet (*Genetta victoriae*) may be eaten only by the elders. Women do not eat this animal because of its strong smell; men cannot eat it otherwise their offspring might have diseases of the respiratory system. The parents of little children cannot eat the spotted-necked otter (*Lutra maculicollis*) because their babies would risk a dangerous case of dysentery. According to the Mbuti Efe (E. Ituri), *Nandinia binotata* and *Civettictis civetta* may not be eaten by pregnant women or by their husbands, otherwise the baby might be born with some anomalies.

All carnivore species may also be totemic animals of single clans for both pygmies and villagers. In that case, they believe these animals are their own ancestors and must never be eaten.

All these food restrictions, together with the good status of forest conservation and low human population density in the Ituri region (3 persons per km<sup>2</sup>) lead us to think that the forest species of Mustelidae and Viverridae will still survive unthreatened for many years. In fact, there is no scientific information on the status of their populations but we can suggest it is satisfactory: most of the species are widespread and common, and perhaps they may tolerate the current hunting pressure. The rarest and endemic species, such as the aquatic genet, *Osbornictis piscivora*, are very elusive and represent only exceptional game for both pygmies and villagers. In the near future, however, deforestation associated with increasing human population and fire-arms introduction will reduce the population size of all the larger mammals to risk levels.

Conservation measures should be taken considering villager demography and livestock absence throughout the Ituri Forest. The increasing Bantu populations of the region survive by the exploitation of game and forest clearing. A large protected area of forest reserve (1.383.000 ha) in the Epulu River region was recently accepted by the government as a potential National Park.

## Virunga National park

The Virunga National Park covers an area of 780.000 ha, within one of the most populated regions of Zaire. More than 14.000 persons live in the Savannah just around Lake Idi Amin; almost one million people inhabit the area surrounding the southern sector of the park (volcanoes and montane forest). Most of the people belong to the Nande and Bwisha tribes of agriculturalists. All these people need firewood and timber for domestic use. Furthermore, many of them set out traps either in the forest or in the savannah to catch small and medium-sized mammals. In the Mitumba Mountains (west of Lake Amin), forests have almost completely disappeared and most of the mammal species are locally extinct.

All of the Mustelidae and Viverridae species are eaten by the Nande, except for the white-naped weasel (*Poecilogale albinucha*) and the striped polecat (*Ictonyx striata*), because of their bad smell. The former is used as a medicine against rheumatic pains: its ashes are put into a small incision of the patient's skin. Genet skins are used in the same way, while the excrements of *Civettictis civetta*

are considered a remedy against infectious diseases.

Genets are frequently poached in the volcanoes forests and many skins are put on sale every day in the tourist shops of Goma, the regional capital, together with those of servals, dassies, and threatened monkeys (golden monkey and black & white colobus).

The most threatened populations of Mustelidae and Viverridae are surely those of the southern sector of the park (montane forests and volcanoes) and the Mitumba Mountains. Luckily, no endemic species seems to occur in these endangered ecosystems. Further taxonomic studies could reveal the presence of endemic subspecies in these yet unknown carnivore populations.

Department of Animal and Human Biology,  
University of Rome I, Italy.

SPECIES	DISTRIBUTION			
<b>Mustelidae</b>			VI	GA
<i>Aonyx capensis</i> (Schinz, 1821)		IF(!)		
<i>Aonyx congica</i> (Lonnberg, 1910)	TM		VI	GA
<i>Lutra maculicollis</i> Lichtenstein, 1835	TM		VI(!)	GA(!)
<i>Mellivora capensis</i> (Schreber, 1776)	TM		VI(!)	
<i>Poecilogale albinucha</i> (Gray, 1864)			VI	
<i>Ictonyx striata</i> (Perry, 1810)				
<b>Viverridae</b>				
<i>Nandinia binotata</i> (Gray, 1830)	TM	IF(!)	VI(!)	
<i>Civettictis civetta</i> (Schreber, 1776)	TM	IF(!)	VI(!)	GA(!)
<i>Osbornictis piscivora</i> J. A. Allen, 1919	TM	IF		
<i>Poiana richardsoni</i> (Thompson, 1842)	TM	IF(!)		
<i>Genetta victoriae</i> Thomas, 1901	TM	IF(!)		
<i>Genetta servalina</i> (Pucheran, 1855)	TM	IF(!)	VI	
<i>Genetta rubiginosa</i> Pucheran, 1855	TM		VI	GA
<i>Genetta pardina</i> (I. Geoffroy, 1832)	TM			
<i>Bdeogale nigripes</i> Pucheran, 1855	TM	IF(!)	VI(!)	
<i>Ichneumia albicauda</i> (G. Cuvier, 1829)			VI	GA(!)
<i>Herpestes ichneumon</i> Linnaeus, 1758	TM		VI(!)	GA(!)
<i>Herpestes sanguineus</i> (Rüppel, 1835)			VI	GA(!)
<i>Herpestes naso</i> (De Winton, 1901)	TM			
<i>Atilax paludinosus</i> (G. Cuvier, 1829)	TM		VI	GA
<i>Mungos mungo</i> (Gmelin, 1788)			VI	GA(!)
<i>Crossarchus alexandri</i> Thomas & Wroughton, 1907	TM	IF(!)	VI	
<i>Dologale dybowskii</i> (Pousargues, 1893)				GA

TM = Tshopo-Maiko Rivers region

IF = Ituri Forest

VI = Virunga National Park

GA = Garamba National Park

# Conservation and economic importance of the mustelids and viverrids in China.

TAN Bangjie

## Status of conservation

Of the 19 species of mustelids only two, the wolverine (*Gulo gulo*) and the sable (*Martes zibellina*), belong nominally to the Chinese Protected Animal List Category I, and five: the stone marten (*Martes foina*), the yellow-throated marten (*Martes flavigula*), the European otter (*Lutra lutra*), the smooth-coated otter (*Lutrogale perspicillata*), and the oriental small-clawed otter (*Amblonyx cinerea*) belong nominally to category II.

The remaining ones, including all the weasels and badgers, are left "unprotected" for the time being.

Of the 11 species of viverrids only one, the binturong (*Arctictis binturong*) is listed in category I, and three others: the large Indian civet (*Viverra zibetha*), the small Indian civet (*Viverricula indica*), and the common palm civet (*Paradoxurus hermaphroditus*) are listed in category II.

The "Chinese Wildlife Conservation Law" adopted by the National People's Congress in November 1988 has become effective from March 1, 1989. According to the law animals of category I are not allowed to be collected without a special permit from the central government authorities, while the collecting of animals of category II needs a permit from the provincial authorities. Various terms of punishment are provided by the law for violators. It is hoped that with the legislation of the law free hunting and poaching may be checked to some extent hereafter.

As for the present status of these animals it seems that none of them is really very endangered. But, speaking as a whole, their present populations are markedly reduced, having been hunted without restriction for so many years.

## Economic value

To what extent the protective regulations may be effective, we will have to wait and see. For generations up to the present, almost all the species mentioned above have been hunted or poached by the masses and collected by commercial agents as fur animals. A few of them have additional economic value: thus both the large and the small Indian civet are "civet" producers, and the masked palm civet (*Paguma larvata*) is considered as a delicacy. Besides being freely on sale in Guangdong and Guangxi markets, masked palm civets (known as "guo-zi-li" or "fruit cat") are exported in thousands to Hong Kong annually for the local gourmet's consumption.

On the whole the economic value of all wild animals is exploited to the utmost by the Chinese. For instance the pelts of the large and small Indian civets, the masked palm civet, the crab-eating mongoose (*Herpestes urva*), and the ferret badgers (*Melogale* sp.) have little value in other countries but are important fur bearers in China. Coats made from such furs are widely on sale on the markets.

The pelts of Eurasian badgers (*Meles meles*) and hog

badgers (*Arctonyx collaris*), being harsh in quality, are well suited for making rugs. In addition their guard hairs are used to make brushes and their fat for medicines.

Most of the Chinese fur animals, regardless of whether their furs are valuable or not, have really little economic significance because the annual harvest is quite negligible. The annual collection of otters, sables, stone martens, and ermines, despite the high cost of their fur, amounts only to some hundreds of each. This is even more the case with the wolverine (*Gulo gulo*), whose pelt is valuable, but whose wild population is so small in China that its annual catch seldom exceeds ten specimens.

Exceptions are Siberian weasels (*Mustela sibirica*), ferret badgers (*Melogale moschata*) and alpine weasels (*Mustela altaica*): although individual pelts are sold much cheaper than those of sables and otters, the collective income is much higher because the annual catches are quantitatively much greater. Statistic figures indicate that the annual average catch of Siberian weasels amounts from 2.5 millions (1972-1977) to 3 millions (1978-1980) with a peak of 3.52 millions. The annual catch of ferret badgers varies between 300,000 and 500,000, of alpine weasels between 250,000 and 300,000. These furs and their products are staple exports for China. Moreover these figures are probably incomplete, as specimens caught for private consumption have not been counted.

Formerly fur coats were seldom worn by Chinese women, but, as a result of the economic reforms, such coats are seen more frequently in Chinese cities lately. A coat made of Siberian weasel- or ferret badger pelts is generally sold for 300-500 US\$ in Chinese fur stores. Coats made from pelts of large Indian-, small Indian civet, common palm-, or masked palm civet are 15 to 30% cheaper than those made of weasel or ferret badger, and also cheaper than those made of leopard cat or marmot.

On the other hand, prices of fur coats from sable, marten or otter are so high, these products are only for export and seldom if ever seen on the local markets.

The tail-hairs of the Siberian weasel and small Indian civet are also used in the production of fine brushes for Chinese painting and writing.

## Distribution

Some species are widely distributed in China. For instance the European otter, the yellow-throated marten, and the two species of badgers, which are found both in the northern and southern provinces. On the other hand most of the martens and weasels are distributed in the north, while the ferret-badgers and the two remaining species of otters live in the south. Most of the mustelids have a fairly extensive range except the wolverine, whose small population is limited to the Greater Xing-an-ling Mountains

in northern Heilongjing Province, and the back-striped weasel (*Mustela strigidorsa*) which is found in Yunnan Province only. The smooth-coated otter (*Lutrogale perspicillata*) is limited to the western part of Yunnan.

Almost all the civets and mongooses are animals of the subtropical regions of south, southwest, and southeast China. The large Indian civet, the small Indian civet, the masked palm civet, and the crab-eating mongoose (*Herpestes urva*) are wide-ranged species with large populations. On the other hand, the binturong (*Arctictis binturong*) is strictly limited to the Xishuangbanna region of Yunnan. Whether or not Lowe's otter civet occurs in China is not yet known. It is worth getting deeper into the distribution of the masked palm civet. It seems that it is the only viverrid whose range of distribution extends from the southern subtropical provinces into the northern temperate provinces. Surveys made in recent years have proved that it is not only found in the southern part of Henan and Shaanxi provinces south of the Yellow River, but also in central Hebei and Shanxi far north of the Yellow River bank, and even in the mountains to the north and west of Beijing up to latitude 40°N. The Beijing Zoo in the fifties received quite a number of specimens snared in the western Hills in the suburbs of Beijing, but no more since the seventies. These specimens were obviously larger, longer haired, and greyer coloured than specimens from the south, so that they may form a different race. As for its population, past records indicated that the annual catch of this species was 80,000 to 100,000 in the 1960's. The present figure is not known. As an unprotected animal it is still freely hunted both for its fur

(known as "qing-zong" or blue-haired in the fur trade) and its meat. The reduction of its population is quite anticipated.

## Farming

Since natural resources are getting more and more scarce and the animals more difficult to collect, people are inclined to raise some of the species with economic value in farms. Experiments in the past dozen years have shown that some are more or less successful while others are not. For instance, a number of farms or experimental stations in northeast China trying to raise sable and Siberian weasel for a profit all ended in failure for one reason or another. Experiments with ferret badgers and marbled polecats (*Vormela peregusna*) are still being continued on a very small scale. As far as mustelids are concerned, the only successful project in China is mink farming. Numerous large and small mink farms are scattered almost all over China. But the animal is not Chinese, the breeding stock being imported from abroad, and its products are for export to foreign countries. The furs and fur coats are too costly to be sold locally.

At the same time, fur farming with canids (foxes and racoon dogs) is also very popular in China.

Meanwhile civet farming has begun to thrive in China. "Civet musk" is accepted as a good supplement to "deer musk", which is very costly and difficult to get. Among the limited civet farms in China, the Hangzhou Zoo is a forerunner. Hundreds of civets of both species are presently being kept by the zoo for musk collection.

### List of the 19 species of mustelids and 11 species of viverrids known to occur in China.

#### A. MUSTELIDS

1. *Gulo gulo* (Linnaeus, 1758) - wolverine
2. *Martes zibellina* (Linnaeus, 1758) - sable
3. *Martes foina* (Erxleben, 1777) - stone marten
4. *Martes flavigula* (Boddaert, 1758) - yellow-throated marten
5. *Mustela erminea* Linnaeus, 1758 - ermine
6. *Mustela sibirica* Pallas, 1773 - Siberian weasel
7. *Mustela altaica* Pallas, 1811 - alpine weasel
8. *Mustela kathiah* Hodgson, 1835 - yellow-bellied weasel
9. *Mustela strigidorsa* Gray, 1853 - back-striped weasel
10. *Mustela putorius* Linnaeus, 1758 - polecat
11. *Mustela nivalis* Linnaeus, 1766 - weasel
12. *Vormela peregusna* (Güldenstaedt, 1770) - marbled polecat
13. *Melogale moschata* (Gray, 1831) - Chinese ferret badger
14. *Melogale personata* Geoffroy, 1831 - Burmese ferret badger
15. *Meles meles* (Linnaeus, 1758) - Eurasian badger
16. *Arctonyx collaris* F. Cuvier, 1825 - hog badger
17. *Lutra lutra* (Linnaeus, 1758) - European otter
18. *Lutrogale perspicillata* (Geoffroy, 1826) - smooth-coated otter
19. *Amblyonyx cinerea* (Illiger, 1815) - oriental small-clawed otter

#### B. VIVERRIDS

1. *Viverra zibetha* Linnaeus, 1758 - large Indian civet
2. *Viverra megaspila* Blyth, 1862 - large-spotted civet
3. *Viverricula indica* (Desmarest, 1817) - small Indian civet
4. *Prionodon pardicolor* Hodgson, 1842 - spotted linsang
5. *Paguma larvata* (Hamilton-Smith, 1827) - masked palm civet
6. *Paradoxurus hermaphroditus* (Pallas, 1777) - common palm civet
7. *Arctogalidia trivirgata* (Gray, 1832) - small-toothed palm civet
8. *Arctictis binturong* (Raffles, 1821) - binturong
9. *Chrotogale owstoni* Thomas, 1912 - Owston's palm civet
10. *Herpestes auropunctatus* (Hodgson, 1836) - small Indian mongoose
11. *Herpestes urva* (Hodgson, 1836) - crab-eating mongoose

Chinese Association of Zoological Gardens, 137  
Xi Zhi Men Wai Street, Beijing, China.

# Conservation of the Liberian mongoose.

Mark E. Taylor

The Liberian mongoose (*Liberiictis kuhni*) was first described in 1958 on the basis of a number of skulls, while the external characteristics of the species were not described until 1974 from two freshly killed animals obtained by Schlitter in 1971.

Although other specimens have been obtained subsequently and used for anatomical research, little ecological information is available for the species. Until 1989 there were no known published photographs of a live animal.

The species is rare and has only ever been seen or collected in Liberia. It is hunted for food by rural Liberians in Nimba and Grand Gedeh Counties, and its ecological requirements are more or less unknown. In preparing the IUCN/SSC Action Plan for Global Mustelid and Viverrid Conservation, the committee recommended that a study of the distribution of the Liberian mongoose should be started together with a captive breeding programme. I undertook to try and accomplish this and in 1988 led my first expedition from the Royal Ontario Museum and Metro Toronto Zoo to Liberia. I found one Liberian mongoose which had been shot by a hunter in the Gbi National Forest near Tappita, Nimba County. Attempts to trap this species in Sapo National Park (Sinoe County) which is some 60 km south of where Schlitter collected his specimens were unsuccessful, though marsh mongooses and a palm civet were caught and kusimanses were seen.

The purpose of the Metro Toronto Zoo 1989 expedition to Liberia was to survey the distribution, abundance and ecological requirements of the Liberian mongoose in Nimba and Grand Gedeh Counties and to obtain if possible two pairs to start a captive breeding program. A large number of Tomahawk live traps of various sizes were used and were set along animal trails, by paths and roads and near watering places in forested and secondary growth areas. They were baited with a variety of foods including fresh meat, fish, crabs and fruits. The trapping rate of viverrids was less than 0.5%, though other vertebrates caught included dwarf crocodile, pigmy rail and giant rats. Local hunters also used our traps but their success rate was no better.

During the 1989 expedition I distributed several hundred coloured brochures to forestry staff, villagers and hunters. These brochures illustrated the Liberian mongoose, Gambian mongoose, slender mongoose, kusimanse, forest genet and African linsang. Discussions with villagers were aimed at determining the relative abundance of the commoner species and to what extent they were aware of the rare species. Many villagers claimed they knew all six

species which is highly unlikely, while others knew some of the species and were able to describe their habits and where they lived. The general information I gathered was that the Liberian mongoose is uncommon, not present in some of the areas visited, is fierce and is found most commonly by water courses, at least during the dry season. It also digs worms out of rotten palm tree stumps.

A single live Liberian mongoose was obtained from a villager, John Weyu, on February 6th, 1989 in the Gbi National Forest. He had received one of our coloured brochures during a previous visit to his village and caught this animal, a young adult male, in a snare. This resulted in damage to the foot and as soon as we obtained the mongoose we started medical treatment with injections of penicillin and surface cleansing with a dilute Betadine solution. The foot responded well to this treatment, and the animal gained weight, and was soon able to walk and trot. He was flown to Canada on March 19, 1989, and is now at the Metro Toronto Zoo.

I hired John Weyu to try and catch another Liberian mongoose, and gave him a number of my Tomahawk traps. In spite of two weeks trapping he was not successful, although he did catch a single kusimanse. The use of these live traps has generally been unsuccessful in Liberia.

I used a remote infra-red beam activated camera which was very useful in surveying animals in the forest. With this equipment I obtained photographs of leopard, civet and genet, and with more time and equipment I am sure that it would be an effective way of censusing an area for terrestrial species. Baiting an area with either meat or fruit assists in attracting species which avoided the live traps.

During the expedition many Liberians were interviewed and the goals of the expedition discussed. The value of animals as an integral and important part of Liberia's heritage was stressed and the uniqueness of the Liberian mongoose was emphasised.

Although I was unsuccessful in obtaining more than a single live animal, I am now familiar with the habitat requirements of this species and have recommended to Wildlife officials of the Liberian Forestry Development Authority that hunting restrictions be enforced in the Gbi National Forest.

160 Woodycrest Avenue  
Toronto  
Ontario M4J 3B9,  
Canada.

# Diversity and conservation status of mustelids in Mexico

GERARDO CEBALLOS

The diversity of native mustelid species in Mexico is relatively high. The following 15 species are distributed in the country: tayra (*Eira barbara*), grison (*Galictis vittata*), striped skunk (*Mephitis mephitis*), hooded skunk (*Mephitis macroura*), three hog-nosed skunks (*Conepatus mesoleucus*, *C. leuconotus*, and *C. semistriatus*), three spotted skunks (*Spilogale gracilis*, *S. putorius*, and *S. pygmaea*), northern river otter (*Lutra canadensis*), southern river otter (*Lutra longicaudis*), sea otter (*Enhydra lutris*), American badger (*Taxidea taxus*), and long-tailed weasel (*Mustela frenata*). The largest species is the tayra (5 kg) and the smallest one the pygmy skunk (0.3 kg) (Leopold, 1959; Hall, 1981; Ceballos & Navarro, in press).

Practically all these mustelid species, but the northern river otter and the sea otter, were widely distributed in Mexico in historic times. Human activities such as indiscriminate hunting and habitat destruction have caused population declines in all the species in recent decades. Most species however, are considered out of danger because they have been able to cope with the transformation of their habitats. Unfortunately, several species are in danger of extinction. Overexploitation of both the sea otter and the northern river otter caused their extirpation from the country. The pygmy skunk, the southern river otter and the grey-headed tayra (*E. b. senex*) are considered threatened or endangered throughout their geographic ranges. The problems faced by the grey-headed tayra and the pygmy spotted skunk are discussed in detail in the Action Plan for the Conservation of Mustelids and Viverrids (Schreiber et al., 1989).

Two species, the American badger and the grison, although not globally endangered, face severe conservation

problems in Mexico (Ceballos & Navarro, in press). The badger populations, who are considered predators of small domestic animals, have been decimated by hunting and poisoning. Predator control campaigns using non-specific pesticides, such as compound 1080 (Sodium fluoracetate), were carried out until recent years. Presently, the use of 1080 is illegal but there are enormous problems to enforce the law. Badgers are protected in La Michilia and Mapimi biosphere reserves in the state of Durango. Badgers are quite abundant in the USA states bordering Mexico.

The conservation status of the grison throughout its geographic range is unknown. There are few records of grison in Mexico, most from tropical rain forests. Such forests are disappearing really fast in the country threatening the long-term survival of this species (Ceballos & Navarro, in press). A few populations are protected in reserves such as Sian Ka'an (Quintana Roo), Los Tuxtlas (Veracruz), and Montes Azules (Chiapas). However, a long-term study of its distribution and ecology is required.

## References

- Ceballos, G. & Navarro, D. In press. Diversity and conservation of Mexican mammals. In: Latin American mammalogy: Ecology, evolution and conservation (M. A. Mares & D. J. Schmidly, eds.). Oklahoma University Press.
- Hall, E. R. 1981. The mammals of North America. Wiley-Interscience, New York.
- Leopold, A. S. 1959. Fauna silvestre de Mexico. IMRNAR, Mexico, D. F.
- Schreiber, A., Wirth, R., Riffel, M. & Van Rompaey, H. 1989. Weasels, civets, mongooses, and their relatives. An Action Plan for the conservation of mustelids and viverrids. IUCN, Gland.

## News from CITES

The Government of India has submitted to the CITES Secretariat the following list of mustelid and viverrid species for inclusion in Appendix III:

<i>Martes flavigula</i>	Yellow-throate marten
(including <i>Martes gwatkinsi</i> , the Nilgiri marten)	
<i>Martes foina intermedia</i>	Central Asian stone marten
<i>Mustela altaica</i>	Mountain weasel
<i>Mustela erminea</i>	Stoat
<i>Mustela kathiah</i>	Yellow-bellied weasel
<i>Mustela sibirica</i>	Siberian weasel
<i>Arctictis binturong</i>	Binturong
<i>Paguma larvata</i>	Masked palm civet
<i>Paradoxurus hermaphroditus</i>	Common palm civet
<i>Paradoxurus jerdoni</i>	Brown palm civet

<i>Viverra megaspila</i>	Large-spotted civet
<i>Viverra zibetha</i>	Large Indian civet
<i>Viverricula indica</i>	Small Indian civet
<i>Herpestes auropunctatus</i>	Small Indian mongoose
<i>Herpestes edwardsi</i>	Indian grey mongoose
<i>Herpestes fuscus</i>	Indian brown mongoose
<i>Herpestes smithii</i>	Ruddy mongoose
<i>Herpestes urva</i>	Crab-eating mongoose
<i>Herpestes vitticollis</i>	Stripe-necked mongoose

The listing of these species became effective on 16 March 1989.

Traffic Bulletin 10(3/4):29.

# Mustelid and viverrid genetics: A call for cooperative action!

ARND SCHREIBER

Imagine that the seafarers who roasted the last dodo had spared a sample of its tissue! We now could extract genomic DNA from it, and carry out a great number of scientific investigations, yielding data on various genetical and physiological details of this peculiar creature, its taxonomy, and its evolutionary relationships. Most of this must remain mysterious because no one cared to conserve some tissue. Still, similar opportunities are lost frequently, in each case when a rarely kept mustelid and viverrid dies, and the keeper does not care to preserve its genome (although this is fairly easy). DNA is a stable molecule contained in any organ and tissue, and it is easily secured if the sample is removed soon, and stored frozen until the nucleic acid is extracted in the laboratory. Once the DNA is solved in an appropriate buffer solution, or lyophilized, it can be stored for many years.

Therefore I ask every interested zoo curator, or private keeper, to cooperate in an effort to collect tissue samples of fresh carcasses of rare mustelids and viverrids for a DNA bank. Inadvertently trapped animals are another possible source (but scientific enthusiasm should not lead trappers to persecute rare species). Collected over a number of years, enough species could be available for comparative purposes, and hopefully, also several individuals from single species for population genetical analysis. There are numerous additional biological fields which could profit from certain molecular analyses, such as conservation genetics, immunology, systematics, evolutionary biology and ecology.

## Background

Hardly any mustelid or viverrid species considered as threatened (or likely to be so) by the Mustelid and Viverrid Specialist Group has been the object of intensive molecular genetical studies, and DNA investigations appear to be totally lacking. The very few genetic studies published about wild mustelids focused on starch gel electrophoresis of selected proteins. I am not aware of a comprehensive genetic study of viverrids using molecular biological methods. On the other hand, carnivore genetics revealed major surprises, including a remarkably low degree of genetic variation in some species. Likewise, little variation was found in two of three recent studies in mustelids.

Hardly anything is known about the immunogenetics of rare mustelids and viverrids. Endemic diseases, such as distemper and rabies, are common in a number of rare mustelid/viverrids populations. The last known population of the black-footed ferret was almost exterminated by canine distemper. While some molecular investigations based on the only few specimens necessarily available in rare species must remain fragmentary for a long time to come, even single data can give hints to understand more effectively the biology of a species. Wouldn't it be a sad waste if the tissues of the only otter civet presently in captivity were not preserved?

## Action

While many keepers will not like this idea while their pet is still living, a small tissue sample will -if treated correctly- yield enough DNA to keep numerous researchers busy. I hope that many keepers will wish to cooperate by preserving at least a whale-nut-sized piece of muscle, and possibly the heart, kidneys, and liver (or anything else needed). Only three aspects must be kept in mind: first, freeze the material as soon as possible after death; second, store it as cold as possible, preferably at  $-80\text{ C}^{\circ}$  (after shock-freezing in liquid nitrogen, or in a mixture of dry ice and alcohol), although storage in an ordinary kitchen freezer reaching  $-20\text{ C}^{\circ}$  is preferable to losing the valuable material; third, make sure that it does not thaw again. Many collections will send their carcasses to veterinary institutes for autopsy, or will wish to mount the specimens for exhibition in a museum. However, samples of soft tissues can usually be taken without interfering with other post-mortem uses. It might be wise to save reasonably sized samples, in view of unforeseen future needs, including those of your own institution. If you agree to cooperate with me, I am certainly willing to store any remaining DNA, and protein fractions, for future uses by you, or interested colleagues. In lyophilized condition, DNA can reach you at each corner of the globe, by air mail, landrover, or on horseback.

The only major effort is to send frozen material to a laboratory, e. g. to the address given below. Certainly, the last Chinese purple-spotted flying mongoose would merit any such effort, and in such urgent cases I could probably arrange the transport. However, my research funds are definitely not sufficient to air-ship two dozen frozen European badgers from an Australian zoo to Heidelberg.

## Request for cooperation

If you agree that our Specialist Group should initiate such a programme, if you are even preparing similar studies and are interested in scientific cooperation, or if you just feel that the genes of your specimens should be preserved, please contact me. My personal scientific interest is the degree of intraspecific genetic variation of some immunologically relevant DNA sequences, and molecular evolution. Carcass samples are one approach to such research, but much additional work could be done if blood, or living tissues can be taken. Further technical details had better be discussed by direct correspondence between those interested.

Arnd Schreiber, c/o Prof. G. Sauer,  
Institute of Virus Research, German Cancer  
Research Centre, Im Neuenheimer Feld 280,  
D-6900 Heidelberg, Federal Republic of Germany.

## Mustelid and viverrid wealth in Sikkim

Locked in by snow-capped Himalayan peaks, Sikkim is bordered by Tibet (China), India, Bhutan, and Nepal (see map). The climate is tropical up to 1,050 m, temperate from 1,050 to 3,650 m, and alpine from 3,650 m up to the perpetual snow line at 4,850 m. While much of it is barren, mountainous, and under perpetual snow, Sikkim, smaller than Yellowstone National Park, is also botanically one of Asia's richest areas. About 4,000 flowering plants and ferns -including 400 orchid varieties- flourish here. Although the small carnivore fauna may not be abundant it is certainly varied with three species of mustelids and four of viverrids still present.

Mrs Usha Ganguli-Lachungpa (Project Officer, Fisheries & Wildlife Circle, Govt. of Sikkim) has kindly reported on the two field trips of the Zoological Survey of India made in Sikkim Dec. 1981-Jan. 1982 to the northern and southern part, and Apr. 1982 to the western and again to the southern part. We are happy to learn that following species are still part of Sikkim's amazing fauna:

1. *Martes flavigula*, yellow-throated marten

A pair was sighted on 10 Jan. 1982 in Phodong (N. Sikkim) at ca 1,500 m ASL.

According to Mrs Ganguli-Lachungpa it is still quite common in northern Sikkim beyond ca 2,000 m.

2. *Mustela sibirica*, Siberian weasel

One individual was sighted on 6 Jan. 1982 in Lachen (N. Sikkim) at ca 2,850 m.

Mrs Ganguli-Lachungpa reports the sighting of a young Siberian weasel at the hot spring at Yumthang (ca 3,700 m), also in northern Sikkim.

3. *Lutra lutra*, European otter


Although the otters are the chief concern of a separate Specialist Group we cannot resist reporting the good news that the European otter was commonly seen in large streams near Hee Gyathang (north) and Melli (south) resp. at ca 1,850 and ca 450 m in Jan. 1982.

4. *Viverra zibetha*, large Indian civet

One individual was observed in each of the following localities: 25-27 Dec. 1981, Singhik (ca 1,400 m, N.); 21-26 Apr. 1982, Naya Bazar (ca 400 m, S.); 23-24 Apr. 1982, Yoksum (ca 1,800 m, W.); 25-26 Apr. 1982, Pelling (ca 2,000 m, W.).

5. *Prionodon pardicolor*, spotted linsang

The report on the spotted linsang is based on the sighting of



one individual by Dr. R. K. Ghose in Lema (ca 2,300 m) in Dec. 1981.

6. *Paguma larvata*, masked palm civet

Masked palm civets were sighted several times: one each in Singhik (ca 1,400 m) and Hee Gyathang (ca 1,800 m) -both in the north-, Yoksum (ca 1,800 m) in the west, and a pair in Naya Bazar (ca 400 m) in the south.

Mrs Ganguli-Lachungpa found that *Paguma larvata*, together with *Viverra zibetha*, and *Lutra lutra*, are still common in east, south, and west Sikkim.

7. *Arctictis binturong*, binturong

A pair of binturongs was observed on 4 Jan. 1982 in Menshithang (ca 3,150 m, N.), and one individual 8-9 Jan. 1982 in Hee Gyathang (ca 1,800 m, N.).

The distribution on the left bank of the Teesta River in northern Sikkim is confirmed by Mrs Ganguli-Lachungpa who found binturongs "not common" there.

## Binturong Studbook announcement

Last May the "Regional Studbook of the Binturong (*Arctictis binturong*) in the British Isles" was published. It was compiled by Paul Robinson of Southport Zoo, Southport, UK. Besides some preliminary chapters on taxonomy and distribution, status in the wild, status and history in captivity, reproduction, and diseases, it contains data on the 63

binturongs kept in nine zoos and one private collection in the UK. The genealogy is current through December 1987.

This is to our knowledge the first and only viverrid studbook in existence at present. We hope this useful and much needed work may serve as a model.

# RESEARCH

## ...from R. A. Mead: The reproductive cycle of the wolverine.

Greg Starypan of Northwest Trek Park, Tacoma, Washington invited me to assist them in studying the reproductive cycle of 15 captive wolverines (*Gulo gulo*) owned by Dale Peterson. Others participating in this study are Marc Rector, graduate student at the University of Washington, and Dr. Mike Jones, DVM. We have now pretty well delineated the testis cycle in the male and have monitored testosterone levels. Testicular recrudescence begins in late January and the testes reach maximal size in May. Two out of two males produced good ejaculates in June. Testicular regression begins in late July and appears to be complete by September. The females exhibited increased vaginal cornification from June to early August; however, few of them exceeded 90% cornified epithelial cells in the vaginal lavages for more than a few days and none of the females bred when exposed to males. Plasma progesterone levels remained basal throughout the study suggesting that the wolverine does not ovulate and form corpora lutea unless bred. This study is now about to enter its second year and we hope to obtain breeding this year.

Dr. Rodney A. Mead, Dept. of Biological Sciences,  
University of Idaho, Moscow, Idaho 83843, USA.

## ...from Gerardo Ceballos:

1. The "Asociacion Mexicana de Mastozoologia" (Mexican Association of Mammalogists), a scientific society devoted to the study and conservation of the mammals of Mexico, welcomes international members. For more information please send letter to:

Dr. Gerardo Ceballo President AMMAC  
Apartado Postal 70-419  
Mexico, D. F. 04510 MEXICO

2. I am beginning a study about the ecology of the pygmy skunk (*Spilogale pygmaea*) at the Biological Station of Chamela, located in western Mexico. Anybody interested please get in touch.

Dr. Gerardo Ceballos,  
Apartado Postal 23-D, Toluca,  
Mex. 50120, Mexico

## ...from U. S. Seal: Request for population viability data.

We have prepared POPULATION VIABILITY ANALYSIS (PVA) Data Forms to begin the process of collecting the information necessary to construct population models as a part of the process of evaluating the status and needs for long term conservation of species and populations in the wild and in captivity. We are preparing an action plan for captive carnivore populations and wish to work closely with the specialist groups. We welcome your comment, interest, and participation.

## POPULATION VIABILITY ANALYSIS DATA FORM - MAMMALS

Species:  
Species distribution:  
Study taxon (subspecies):  
Study population location:  
Metapopulation -are there other separate populations?  
Are maps available? Separation by distance, geographic barriers?:  
Specialized requirements (trophic, ecological):  
Age of first reproduction for each sex (proportion breeding):  
a) Earliest:  
b) Mean:  
Gestation period (days or weeks):  
Litter size (N, mean, SD, range) (at birth?, weaning?):  
Birth season:  
Birth frequency (interbirth interval):  
Reproductive life-span (male & female, range):  
Life time reproduction (mean, male & female):  
Adult sex ratio:  
Adult body weight of males and females:  
Social structure in terms of breeding (random, pair-bonded, polygyny, polyandry, etc.; breeding male and female turnover each year?):  
Proportion of adult males and females breeding each year:  
Dispersal distance (mean, sexes):  
Migrations (months):  
Territoriality (home range, season):  
Birth sex ratio:  
Birth weights (male & female):  
Ovulation -induced or spontaneous:  
Implantation -immediate or delayed (duration):  
Estrous cycles (seasonal, multiple or single, post partum):  
Duration of lactation:  
Post-lactational estrus:  
Age of dispersal:  
Maximum longevity:  
Population census -most recent. Date of last census.  
Reliability estimate:  
Projected population (5, 10, 50 years):  
Past population census (5, 10, 20 years -dates, reliability estimates):  
Population sex and age structure (young, juvenile, and adults) -time of year:  
Fecundity rates (by sex and age class):  
Mortality rates and distribution (by sex and age) (neonatal, juvenile, adult):  
Population density estimate. Area of population. Attach marked map:  
Sources of mortality -% (natural, poaching, harvest, accidental, seasonal?):  
Habitat capacity estimate (has capacity changed in past 20, 50 years?):  
Present habitat protection status:  
Projected habitat projection status (5, 10, 50 years):  
Environmental variance affecting reproduction and mortality (rainfall, prey, predators, disease, snow cover?):  
Is pedigree information available?:  
Attach life table if available.  
Data form completed:  
Correspondent/Investigator:

Name:  
Address:  
Telephone:  
Fax:  
References:  
Comments:

U. S. Seal, Chairman IUCN/SSC Captive Breeding Specialist Group,  
12101 Johnny Cake Ridge Road, Apple Valley, MN 55124,  
USA. Tel. (612)431-93245.

### Request for binturong (*Arctictis binturong*) information

Little taxonomic work has been done in recent years on the Asian viverrids. The binturong is no exception. Some members of the M&VSG chose the binturong as study material because, except for some social mongooses, it is the most frequently kept viverrid in zoological collections, and also because a start has been made by Paul Robinson with the publication of a regional studbook.

We would be grateful to hear from any zoo who has kept or is keeping binturongs. We are chiefly interested in

knowing the origins (land, region or locality) of the animal in post mortem findings and we also hope to find a willing participant to part with some tissue samples or blood (see Arnd Schreiber's "Mustelid and viverrid genetics" in this issue for number).

Please contact:  
**Paul Robinson**  
Curator Southport Zoo  
Princes Park,  
Southport, Lancashire, UK

or: **Arnd Schreiber**  
Zoologisches Institut  
Im Neuenheimer Feld, 28  
D-9000 Heidelberg, FGR

or: **Harry Van Rompaey**  
Jan Verbertlei, 15  
2520 Edegem, Belgium

## ABSTRACTS

### American mink in beaver's sites in Poland.

After intentional introduction of American mink (*Mustela vison*) into Byelorussia (USSR) this feral population spread to the west. In the early 1980's the feral minks from the Piska Wilderness (ca. 160 km N. of Warsaw) and those in the Mazurian Lakeland were studied. On those territories only the American mink was found to live. The minks chose beaver sites as their winter home. They were common in beaver burrows, lodges, and food storages; it seems that the feral mink forced the beavers out of their lodges by its permanent presence. Additional research is necessary to find out whether the feral mink is able to reduce the number of beaver kits. Hunters think that the spread of feral mink populations remarkably reduces the number of musk rats and waterfowl.

Zurowski, W. & Kammler, J. 1987. Norka amerykańska (*Mustela vison* Schreber, 1777) w stanowiskach bobrów. *Przegląd Zoologiczny* 31(4):513-521.

### Crab-eating mongoose in Peninsular Malaysia.

D. R. Wells and C. M. Francis report a sighting of a party of three crab-eating mongooses (*Herpestes urva*) in selectively logged (subsequently clear-felled) semi-deciduous, plains-level forest at 6°40'N, 100°14'E, Mata Ayer Forest Reserve, Perlis State in March 1987. Also, the discovery in the collections of the Institute for Medical Research, Kuala Lumpur of an overlooked specimen from the Gombak Valley of Selangor State, at 3°15'N, 101°44'E.

Year of collection (1970) and precise location suggest selectively logged, fully evergreen forest at about 150 m ASL. Together these records expand the southeast Asian range of *H. urva* southward by approximately 500 km. Wells, D. R. & Francis, C. M. 1988. Crab-eating mongoose *Herpestes urva*, a mammal new to Peninsular Malaysia. *Malay. Nat. J.*, 42:37-41.

### Distribution and taxonomy of *Herpestes* sp. in Peninsular Malaysia

Malayan records of *Herpestes brachyurus* and *H. urva* have been exclusively from lowland rain forest, *brachyurus* at low-lying sites north to Perak and Terengganu States, *urva* to approximately 150 m ASL south to Selangor State. External and skull dimensions of individuals of known sex/age class adequately demonstrate that small and large, short-furred, rufous-faced mongooses inhabiting lowland non-forest environments in Malaya (Peninsular Malaysia) south to Selangor, and elsewhere in southeast Asia, constitute a single, sexually dimorphic species, *Herpestes javanicus*. Transfer of the small-sized taxon *perakensis* to *javanicus* plus dismissal of an invalid record eliminates a claimed area of sympatry between this species and *Herpestes auropunctatus*, removing the latter from the Malayan fauna. *Herpestes edwardsi* is probably now extinct in this area.

Wells, D. R. In press. Notes on the distribution and taxonomy of Peninsular Malaysian mongooses (*Herpestes*). *Nat. Hist. Bull. Siam Soc.*, 37(1).

# Recent Literature

## Mustelidae

- Addison, E. M., Strickland, M. A., Stephenson, A. B. & Hoeve, J. 1988. Cranial lesions possibly associated with *Skrjabinylus* (Nematoda: Metastrongyloidae) infections in martens, fishers, and otters. *Can. J. Zool.*, 66(10):2155-2159.
- Anderson, B. F. 1988. Occurrence of the least weasel (*Mustela nivalis*) in Cumberland county, Tennessee. *J. Tenn. Acad. Sci.*, 63(3):83.
- Appel, M. J. G. & Harris, W. V. 1988. Antibody titers in domestic ferret jills and their kits to canine distemper virus vaccine. *J. Amer. Vet. Med. Assoc.*, 193(3):332-333.
- Apps, P. J., Viljoen, H. W., Pretorius, V. & Rohwer, E. R. 1988. Volatile components of the anal gland secretion of the striped polecat *Ictonyx striatus*. *S. Afr. J. Zool.*, 23(2):136-137.
- Arthur, S. M. 1988. An evaluation of techniques for capturing and radiocollaring fishers. *Wildl. Soc. Bull.*, 16(4):417-421.
- Banci, V. & Harestad, A. 1988. Reproduction and natality of wolverine (*Gulo gulo*) in Yukon. *Ann. Zool. Fenn.*, 25(4):265-270.
- Barabasz, B., Jarosz, S. & Kasperczyk, J. 1987. The effect of covering dates on the pregnancy duration, number of litters and fertility in minks. *Zesz. Probl. Postepow. Nauk Roln.*, 341:383-393.
- Berria, M., DeSantis, M. & Mead, R. A. 1988. Effects of supra chiasmatic nuclear ablation and melatonin on delayed implantation in the spotted skunk. *Neuroendocrinology* 48:371-375.
- Berria, M., Joseph, M. M. & Mead, R. A. 1989. Role of prolactin and luteinizing hormone in regulating timing of implantation in the spotted skunk. *Biol. Reprod.*, 40:232-238.
- Biggings, D. E. & Schroeder, M. H. 1988. Historical and present status of the black-footed ferret. *U. S. For. Serv. Gen. Tech. Rep.*, RM-154:93-97.
- Black, R. A. R. 1988. Some observations on a captive honey badger (*Mellivora capensis*). *Naturalist (S. Afr.)*, 32(2):14-18.
- Bock, W. F. 1988. (Investigations on the sites and microclimate of badger setts (*Meles meles*)). *Z. Jagdwiss.*, 34(3):141-152. In German, English summ.
- Bock, W. F. 1988. (The importance of soil composition for the size of badger setts (*Meles meles*), in two areas in S. E. Germany). *Z. Säugetierk.*, 53(6):349-357. In German, English summ.
- Boissin-Agasse, L., Jacquet, J. M., Lacroix, A. & Boissin, J. 1988. Long-term effects of pinealectomy on testicular function, LHRH hypothalamic system, and plasma prolactin levels in the mink, a short-day breeder. *J. Pineal. Res.*, 5(4):385-396.
- Borvinkin, A. D., Chernov, S. M. & Gribanova, L. Y. 1987. Detection of rabies virus in animal brain and salivary glands, using the immunoenzyme method. *Vopr. Virusol.*, 32(6):747-750.
- Briggs, D. J., Briggs, J. M. & Howard, D. R. 1988. Prevalence of rabies in the striped skunk (*Mephitis mephitis*) in Kansas from 1966 to 1986. *Trans. Kans. Acad. Sci.*, 91(3/4):123-131.
- Buskirk, S. W., Harlow, H. J. & Forrest, S. C. 1988. Temperature regulation in American marten (*Martes americana*) in winter. *Natl. Geogr. Res.*, 4(2):208-218.
- Buskirk, S. W. & Harlow, H. J. 1989. Body-fat dynamics of the American marten (*Martes americana*) in winter. *J. Mammal.*, 70(1):191-193.
- Buskirk, S. W. & Lindstedt, S. L. 1989. Sex biases in trapped samples of Mustelidae. *J. Mammal.*, 70(1):88-97.
- Buskirk, S. W., Forrest, S. C., Raphael, M. G. & Harlow, H. J. 1989. Winter resting site ecology of marten in the central Rocky Mountains. *J. Wildl. Manage.*, 53(1):191-196.
- Campbell, T. M. III, Clark, T. W., Richardson, L., Forrest, S. C. & Houston, B. R. 1987. Food habits of Wyoming black-footed ferrets. *Am. Midl. Nat.*, 117(1):208-210.
- Cheeseman, C. L., Cresswell, W. J., Harris, S. & Mallinson, P. J. 1988. Comparison of dispersal and other movements in two badger (*Meles meles*) populations. *Mammal Rev.*, 18(1):51-59.
- Cheeseman, C. L., Wilesmith, J. W., Stuart, F. A. & Mallinson, P. J. 1988. Dynamics of tuberculosis in a naturally infected badger population. *Mammal Rev.*, 18(1):61-72.
- Choate, J. R., Langley, W. & Bailey, V. 1988. The least weasel in southeastern Kansas. *Prairie Nat.*, 20(1):57.
- Clapperton, B. K., Minot, E. O. & Crump, D. R. 1989. Scent lures from anal sac secretions of the ferret *Mustela furo* L. *J. Chem. Ecol.*, 15(1):291-308.
- Clark, B. S. & Clark, B. K. 1988. First record of the least weasel in Oklahoma. *Prairie Nat.*, 20(3):134.
- Clark, T. W. 1987. Restoring balance between the endangered black-footed ferret (*Mustela nigripes*) and human use in the Great Plains and intermountain west. *J. Wash. Acad. Sci.*, 77(4):168-173.
- Clark, T. W., Grensten, J., Gorges, M., Crete, R. & Gill, J. 1987. Analyses of black-footed ferret translocation sites in Montana. *Prairie Nat.*, 19(1):43-56.
- Clements, E. D., Neal, E. G. & Yalden, D. W. 1988. The national badger sett survey. *Mammal Rev.*, 18(1):1-9.
- Courtney, R. F. 1988. Identifying long-tailed weasels on the prairies. *Blue Jay* 46(2):93-96.
- Cresswell, W. J. & Harris, S. 1988. The effects of weather conditions on the movements and activity of badgers (*Meles meles*) in a suburban environment. *J. Zool. (London)* 216(1):187-194.
- Cresswell, W. J. & Harris, S. 1988. Foraging behaviour and home-range utilization in a suburban badger (*Meles meles*) population. *Mammal Rev.*, 18(1):37-49.
- David, P. G. 1988. Further distribution of *Mustela nivalis* in Kentucky. *Trans. Ky. Acad. Sci.*, 49(1-2):37.
- Davies, J. M., Lachno, D. R. & Roper, T. J. 1988. The anal gland secretion of the European badger (*Meles meles*) and its role in social communication. *J. Zool. (London)* 216(3):455-463.
- Dayan, T. & Tchernov, E. 1988. On the first occurrence of the common weasel (*Mustela nivalis*) in the fossil record of Israel. *Mammalia* 52(2):165-168.

- Delibes, M. & Amores, F. 1986(1988). The stone marten *Martes foina* (Erxleben, 1777) (Mammalia, Carnivora) from Ibiza (Pitiusic, Balearic Islands). *Misc. Zool.*, 10:335-345.
- Dillberger, J. E. & Altman, N. H. 1989. Neoplasia in ferrets: eleven cases with a review. *J. Comp. Pathol.*, 100(2):161-176.
- Ekstrom, J., Hakanson, R., Mansson, B. & Tobin, G. 1988. Tachykinin involvement in parasympathetic nerve-evoked salivation of the ferret. *Br. J. Pharmacol.*, 94(3):707-712.
- Elsheikh, L. E., Karlsson, K.-A., Bergman, R. & Abaas, S. 1988. Induction of protective immune response by vaccination against *Pseudomonas pneumonia* of mink. *J. Vet. Med. Ser. B* 35(4):256-263.
- Erlinge, S. & Sandell, M. 1988. Coexistence of stoat, *Mustela erminea*, and weasel, *M. nivalis*: social dominance, scent communication, and reciprocal distribution. *Oikos* 53(2):242-246.
- Fagerstone, K. A. & Johns, B. E. 1987. Transponders as permanent identification markers for domestic ferrets, black-footed ferrets, and other wildlife. *J. Wildl. Manage.*, 51(2):294-297.
- Forrest, S. C., Biggins, D. E., Richardson, L., Clark, T. W., Campbell, T. M. III, Fagerstone, K. A. & Thome, E. T. 1988. Population attributes for the black-footed ferret (*Mustela nigripes*) at Meeteetse, Wyoming, 1981-1985. *J. Mammal.*, 69(2):261-273.
- Fowler, P. A. & Racey, P. A. 1988. Overwintering strategies of the badger, *Meles meles*, at 57°N. *J. Zool. (London)* 214(4):635-651.
- Fuller, T. K., Johnson, W. E., Franklin, W. L. & Johnson, K. A. 1987. Notes on the Patagonian hog-nosed skunk ( *Conepatus humboldti*) in southern Chile. *J. Mammal.*, 68(4):864-867.
- Gill, J. M., Hartley, W. J. & Hodgkinson, N. L. 1988. An outbreak of post-vaccinal suspected distemper-like encephalitis in farmed ferrets (*Mustela putorius furo*). *N. Z. Vet. J.*, 36(4):173-176.
- Gliwicz, J. 1988. Sexual dimorphism in small mustelids: body diameter limitation. *Oikos* 53(3):411-414.
- Głowacinski, Z. & Profus, P. 1987. Next record of the steppe polecat *Mustela eversmanni* Lesson, 1827, in Poland. *Przegl. Zool.*, 31(4):522-526.
- Golovin, S. Y. et al. 1988. Synthesis, cloning and sequencing of the proopiomelanocortin complementary DNA from the pituitary gland of mink *Mustela vison*. *Biorg. Khim.*, 14(2):273-275.
- Gremillion-Smith, C. & Woolf, A. 1988. Epizootiology of skunk rabies in North America. *J. Wildl. Dis.*, 24(4):620-626.
- Groves, C. R. 1988. Distribution of the wolverine in Idaho as determined by mail questionnaire. *Northwest Sci.*, 62(4):181-195.
- Hancox, M. 1988. Field age determination in the European badger. *Rev. Ecol. Terre Vie* 43(4):399-404.
- Hancox, M. 1988. Pine marten recolonisation in Britain. *Int. Zoo News* 35/4(209):12-13.
- Hancox, M. 1988. Dental anomalies in the Eurasian badger. *J. Zool.*, 216(4):606-608.
- Hartl, G. B., Willing, R., Grillitsch, M. & Kiansek, E. 1988. Biochemical variation in Mustelidae: Are carnivores genetically less variable than other mammals? *Zool. Anz.*, 221(1/2):81-90.
- Heggenes, J. & Borgstrom, R. 1988. Effect of mink, *Mustela vison* Schreber, predation on cohorts of juvenile atlantic salmon, *Salmo salar* L., and brown trout, *S. trutta* L., in three small streams. *J. Fish Biol.*, 33(6):885-894.
- Heran, I. 1984. Comments on the age-conditioned changes of skull proportions in some species of Mustelidae. *Lynx* (Prague), 22:11-14. In Czech.
- Heran, I. & Sladek, J. 1984. Bibliography of the Czech research in Carnivora 1970 to 1980. *Lynx* (Prague), 22:121-144. In Czech.
- Hofer, H. 1988. Variation in resource presence, utilization and reproductive success within a population of European badgers (*Meles meles*). *Mammal Rev.*, 18(1):25-36.
- Hoover, J. P., Baldwin, C. A. & Rupprecht, C. E. 1989. Serologic response of domestic ferrets (*Mustela putorius furo*) to canine distemper and rabies virus vaccines. *J. Am. Vet. Med. Assoc.*, 194(2):234-238.
- Humphrey, T. J. & Bygrave, A. 1988. Abortion in a cow associated with Salmonella infection in badgers. *Vet. Rec.*, 123(6):160.
- Isakova, G. K. 1987. Cytogenetic investigation of American mink (*Mustela vison*) bred in fur farm condition. *Sov. Genet. (Engl. transl. Genetika)* 23(6):764-768.
- Jedrzejewska, B. 1987. Reproduction in weasels *Mustela nivalis* in Poland. *Acta Theriol.*, 32(21-31):493-496.
- Johnson, T. B. 1987. Black-footed ferret. *Wildl. Views*, 30(1):17.
- Joseph, M. M. & Mead, R. A. 1988. Size distribution of ferret luteal cells during pregnancy. *Biol. Reprod.*, 39:1159-1169.
- Khlebodiarova, T. M. et al. 1988. The mink proopiomelanocortin gene: characterization of complementary DNA and chromosomal localization. *Genomics* 2(3):185-188.
- Kirkland, G. L. Jr & Maldonado, J. E. 1988. Patterns of variation in cranial damage attributable to *Skrjabinogylus* sp. (Nematoda, Metastrongyloidea) in skunks (Mammalia, Mustelidae) from Mexico. *Southwest. Nat.*, 33(1):15-20.
- Kondo, N., Saitoh, T., Uruguchi, K. & Abe, H. 1988. Age determination, growth and sexual dimorphism of the feral mink (*Mustela vison*) in Hokkaido. *J. Mammal. Soc. Jpn.*, 13(1):69-75.
- Korhonen, H. 1988. Seasonal comparison of body composition and hair coat structure between mink and polecat. *Comp. Biochem. Physiol. A Comp. Physiol.*, 91(3):469-473.
- Korhonen, H., Harri, M. & Mononen, J. 1989. Regulation of weight loss in male farm mink. *Comp. Biochem. Physiol. A Comp. Physiol.*, 92(3):355-357.
- Krott, P. 1985. Weitere Bemerkungen über das Steinmardervorkommen (*Martes foina* Erxleben, 1777) in Kleinsolkta (Mammalia, Carnivora). *Mitt. Abl. Zool. Landesmus. Joanneum*, 33:7-12.
- Kvam, T., Overskaug, K. & Sorensen, O. J. 1988. The wolverine *Gulo gulo* in Norway. *Lutra* 31(1):7-20.
- Laing, R. 1987. The reestablishment of the black-footed ferret to the Canadian prairie. *Prov. Mus. Alberta Nat. Hist. Occas. Pap.*, 9:293-297.
- Lamy, E. & Canivenc, R. 1988. (Photoperiodic reactivation of corpora lutea formed by gonadotrophic hormones in European badger *Meles meles* L.) *C. R. Acad. Sci. Ser. III Sci. Vie* 307(15):759-762. In French.
- Lawes, I. N. C. & Andrews, P. L. R. 1987. Variation of the ferret skull (*Mustela putorius furo* L.) in relation to stereotaxic landmarks. *J. Anat.*, 154:157-171.
- Lobachev, Y. S. & Baidavletov. 1986. *Martes martes uralensis* in the Ural Basin. Pp. 48-49 in: Gvozdev, E. V. (ed.) *Rare species of animals in Kazakhstan*. Nauka, Alma Ata, Kazakhstan, 246 pp. In Russian.
- Lups, P., & Roper, T. J. 1988. Tooth size in the European badger (*Meles meles*) with special reference to sexual dimorphism, diet and intraspecific aggression. *Acta Theriol.*, 33(2):21-33.

- Lups, P., Roper, T. J. & Stocker, G. 1987. Stomach contents of badgers (*Meles meles* L.) in central Switzerland. *Mammalia* 51(4):559-570.
- Magoun, A. J. 1987. Summer and winter diets of wolverines, *Gulo gulo*, in Arctic Alaska. *Can. Field-Nat.*, 101(3):392-397.
- Maguire, L.A., Clark, T. W., Crete, R., Cada, J., Groves, C., Shaffer, M. L. & Seal, U. S. 1988. Black-footed ferret recovery in Montana: a decision analysis. *Wildl. Soc. Bull.*, 16(2):111-120.
- Matyushkin, E. N. 1987. Hunting the musk deer by the yellow-throated marten in the Sikhote-Alin Mountains. *Byull. Mosk. O-va. Ispyt. Prir. Otd. Biol.*, 92(5):28-42. In Russian.
- Mead, R. A. & Neirinckx, S. 1989. Hormonal induction of oestrus and pregnancy in anoestrous ferrets (*Mustela putorius furo*). *J. Reprod. Fert.*, 86:in press.
- Mead, R. A., Joseph, M. M., Neirinckx, S. 1988. Optimal dose of human chorionic gonadotropin for inducing ovulation in the ferret. *Zoo Biol.*, 7:263-267.
- Mead, R. M., Joseph, M. M., Neirinckx, S. & Berria, M. 1988. Partial characterization of a luteal factor that induces implantation in the ferret. *Biol. Reprod.*, 38:798-803.
- Miller, B. J. & Anderson, S. H. 1989. Failure of fertilization following abbreviated copulation in the ferret (*Mustela putorius furo*). *J. Exp. Zool.*, 249(1):85-89.
- Miller, B. J., Menkens, G. E. & Anderson, S. H. 1988. A field habitat model for black-footed ferrets. *U. S. For. Serv. Gen. Tech. Rep.* RM-154:98-102.
- Miller, B. J., Anderson, S. H., DonCarlos, M. W. & Thorne, E. T. 1988. Biology of the endangered black-footed ferret and the role of captive propagation in its conservation. *Can. J. Zool.*, 66:765-773.
- Moore, D. R. & Kowalchuk, N. E. 1988. Auditory brainstem of the ferret: effects of unilateral cochlear lesions on cochlear nucleus volume and projections to the inferior colliculus. *J. Comp. Neurol.*, 272(4):503-515.
- Moreno, S., Rodriguez, A. & Delibes, M. 1988. Summer foods of the pine marten (*Martes martes*) in Majorca and Minorca, Balearic Islands. *Mammalia* 52(2):289-291.
- Nagorsen, D. W., Forsberg, J. & Giannico, G. R. 1988. An evaluation of canine radiographs for sexing and aging Pacific coast martens. *Wildl. Soc. Bull.*, 16(4):421-426.
- Neal, E. 1988. The stomach contents of badgers, *Meles meles*. *J. Zool. (London)* 215(2):367-369.
- Oakleaf, B. 1988. Why the Meeteetse ferrets had to come in. *Wyo. Wildl.*, 52(3):28-33.
- Paunovich, R. & Forrest, S. C. 1987. Activity of a wild black-footed ferret litter. *Prairie Nat.*, 19(3):159-162.
- Pavlinov, I. Y. & Rossolimo, O. L. 1987. Systematics of the mammals of USSR. *Sb. Tr. Zool. Muz. Mosk. Univ.*, 25:1-284. In Russian.
- Pigozzi, G. 1988. Diet of the European badger (*Meles meles*) in the Maremma Natural Park, central Italy. *Mammal Rev.*, 18(1):73-74.
- Pigozzi, G. 1988. The capture and immobilization of the European badger, *Meles meles* (L.), in its natural environment. *Atti Soc. Ital. Nat. Mus. Civ. Stor. Nat. Milano* 129(1):56-70.
- Powers, B. E., Larue, S. M., Withrow, S. J., Straw, R. C. & Richter, S. L. 1988. Jamshidi needle biopsy for diagnosis of bone lesions in small animals. *J. Amer. Vet. Med. Assoc.*, 193(2):205-210.
- Pullianen, E. 1988. Ecology, status and management of the Finnish wolverine *Gulo gulo* populations. *Lutra* 31(1):21-28.
- Pybus, M. J. 1988. Rabies and rabies control in striped skunks (*Mephitis mephitis*) in three prairie regions of western north America. *J. Wildl. Dis.*, 24(3):434-449.
- Pybus, M. J. 1988. Rabies control by skunk depopulation in southern Alberta, 1983 to 1986. *Prairie Nat.*, 20(1):7-20.
- Rehg, J. E., Gigliotti, F. & Stokes, D. C. 1988. Cryptosporidiosis in ferrets. *Lab. Animal. Sci.*, 38(2):155-158.
- Richardson, L., Clark, T. W., Forrest, S. C. & Campbell, T. M. III. 1987. Winter ecology of black-footed ferrets (*Mustela nigripes*) at Meeteetse, Wyoming. *Am. Midl. Nat.*, 117(2):225-239.
- Sadykhov, I. A. & Fataliev, G. G. 1987. Ecological characteristics of the helminth-faunistic complex of the badger in the lesser and greater Caucasus within the Azerbaijan SSR and pathways of its formation. *Izv. Akad. Nauk Az. SSR Ser. Biol. Nauk* 0(2):80-86. In Russian.
- Schmid, P. & Lups, P. 1988. (The importance of wasps as food for badgers (*Meles meles* L.)). *Bonn. Zool. Beitr.*, 39(1):43-47. In German, English summ.
- Sheng Helin. 1987. Sexual dimorphism and geographical variation in the body size of the yellow weasel (*Mustela sibirica*). *Acta Theriol. Sin.*, 7(2):92-95. In Chinese, English summ.
- Skinner, C. A. & Skinner, P. J. 1988. Food of badgers (*Meles meles*) in an arable area of Essex. *J. Zool. (London)* 215(2):360-362.
- Skrede, A. & Nes, I. F. 1988. Slaughterhouse by-products preserved by *Lactobacillus plantarum* fermentation as feed for mink and foxes. *Animal. Feed. Sci.-Technol.*, 20(4):287-289.
- Smal, C. M. 1988. The American mink *Mustela vison* in Ireland. *Mammal Rev.*, 18(4):201-208.
- Smith, G. L. & Allen, D. G. 1988. Effects of metabolic blockade on intracellular calcium concentration in isolated ferret ventricular muscle. *Circ. Res.*, 62(6):1223-1236.
- Songh, J.-H. et al. 1987. The behaviour of mother and baby sables reared in cage. *Chin. J. Zool.*, 22(5):26-28. In Chinese.
- Stenseth, N. C. & Steen, H. 1987. Dynamics of small populations with special reference to bears, wolverines, and wolves in Norway: a theoretical study. *Viltreport*, 45:1-53.
- Stephenson, R., Butler, P. J., Dunstone, N. & Woakes, A. J. 1988. Heart rate and gas exchange in freely diving American mink (*Mustela vison*). *J. Exp. Biol.*, 134:435-442.
- Storch, I. 1988. (Home range utilization by pine martens.) *Z. Jagdwiss.*, 34(2):115-119. In German.
- Stuart, F. A., Mahmood, K. H., Stanford, J. L. & Pritchard, D. G. 1988. Development of diagnostic tests for, and vaccination against, tuberculosis in badgers. *Mammal Rev.*, 18(1):74-75. Abstract only.
- Sundqvist, C., Amador, A. G. & Bartke, A. 1989. Reproduction and fertility in the mink (*Mustela vison*). *J. Reprod. Fert.*, 85(2):413-441.
- Tait, A. R., Du Boulay, M. & Knight, P. R. 1988. Alterations in the course of and histopathologic response to influenza virus infections produced by enfluzane, halothane, diethyl ether anesthesia in ferrets. *Anesth. Analg.*, 67(7):671-676.
- Tang, Y. P. & Sisk, C. L. 1988. differential effects of testosterone, 5-dihydrotestosterone and estradiol-17 $\beta$  on plasma concentration of LH in castrated ferrets. *J. Endocrinol.*, 117(3):461-466.
- Thorne, E. T. & Williams, E. S. 1988. Disease and endangered species: the black-footed ferret as a recent example. *Conserv. Biol.*, 2(1):66-74.
- Thornton, P. S. 1988. Density and distribution of badgers in south-west England - A predictive model. *Mammal Rev.*, 18(1):11-23.

Timmermans, G. 1988. (First record of the pine marten *Martes martes* west of Amsterdam, The Netherlands.) *Lutra* 31(1):93.

Trapezov, O. V. 1987. Selected transformation of defensive reactions to man in the American mink (*Mustela vison* Schreb.). *Sov. Genet.* (Engl. transl. *Genetika*) 23(6):781-786.

Uraguchi, K., Saitoh, T., Kondo, N. & Abe, H. 1987. Food habits of the feral mink (*Mustela vison* Schreber) in Hokkaido. *J. Mammal. Soc. Jpn.*, 12(1-2):57-67.

Voronov, G. A. 1982. *Acclimatization of mammals on Sakhalin and the Kinil Islands. Results and prospects.* Nauka, Moscow, 133 pp. In Russian.

Weber, D. 1988. (Distribution of polecats (*Mustela putorius* L.) in Switzerland.) *Rev. Suisse Zool.*, 95(4):1041-1056. In German, English summ.

Weber, D. 1989. The ecological significance of resting sites and the seasonal habitat changes in polecats (*Mustela putorius*). *J. Zool.*, 217(4):629-638.

Wiig, O. 1989. Craniometric variation in Norwegian wolverines *Gulo gulo* L. *Zool. J. Linn. Soc.*, 95(3):177-204.

Williams, E. S., Thorne, E. T., Appel, M. J. G. & Belitsky, D. W. 1988. Canine distemper in black-footed ferrets (*Mustela nigripes*) from Wyoming. *J. Wildl. Dis.*, 24(3):385-398.

Wolfel, H. & Schneider, E. 1988. (Studies of set building behaviour of badgers (*Meles meles*)). *Z. Jagdwiss.*, 34(1):47-49. In German, English and French summ.

Yalden, D. W. & Hosey, G. R. (eds.) 1988. Proceedings of a symposium on the badger *Meles meles*, University of Southampton, 28 June 1986. *Mammal Rev.*, 18(1):1-75.

Zheng Shengwi, Li Guihu, Song Shiyang, Han Yiping & Ma Zhaoyun. 1988. Study of ecology of sand badger. *Acta Theriol. Sin.*, 8(1):65-72. In Chinese.

Zielinski, W. J. 1988. The influence of daily variation in foraging cost on the activity of small carnivores. *Anim. Behav.*, 36(1):239-249.

## Mustelidae & Viverridae

Ansell, W. F. H. & Dowsett, R. J. 1988. *Mammals of Malawi. An annotated check list and atlas.* The Trendline Press, Zennor, 170 pp.

Colyn, M. & Van Rompaey, H. 1989. Review of the frontal sinus parasites of Mustelidae and Viverridae, with new data from tropical Africa. *Rev. Zool. Afr.*, 103(1):5-20.

Colyn, M., Dudu, A. & Mankoto ma Mbaelele, M. 1988. Données sur l'exploitation du "petit et moyen gibier" des forêts ombrophiles du Zaïre. 1. Consommation qualitative dans le milieu rural. 2. Commercialisation du gibier en milieu urbain: le cas de Kisangani. *Symp. Int. Conf. "Gestion de la faune en Afrique sub-saharienne"*. Harare, Zimbabwe, 6-12 Oct. 1987, pp. 110-145.

Gao Yaoting et al. 1987. *Fauna Sinica. Mammalia. Vol. 8: Carnivora.* Science Press, Beijing, 377 pp.

Gittleman, J. (ed.) 1989. *Carnivore behaviour, ecology and evolution.* Chapman & Hall, 624 pp.

Schreiber, A., Wirth, R., Riffel, M. & Van Rompaey, H. 1989. *Weasels, civets, mongooses, and their relatives. An action plan for the conservation of mustelids and viverrids.* IUCN, Gland, 100 pp.

Stuart, C. & T. 1988. *Field guide to the mammals of southern Africa.* Struik, Cape Town, 272 pp.

Sumner, J. W., Shaddock, J. H., Wa Guangjer & Baer, G. M. 1987. Oral administration of an attenuated strain of canine adenovirus (type 1) to raccoons, foxes, skunk, and mongoose. *Amer. J. Vet. Res.*, 49(2):169-171.

## Viverridae

Al-Khalili, A. D. 1988. Post-embryonic development of the Indian grey mongoose, *Herpestes edwardsiferrugineus* (Geoffroy, 1818) on the Bahrain Islands. *Afr. Small Mamm. Newsl.*, 10:21-22.

Amr, Z. S., Woodbury, S. C. & Disi, A. M. 1987. On a collection of mammals from Jordan. *Dirasat* 14(12):131-136.

Baker, C. M. 1988. Scent marking behaviour in captive water water mongooses (*Atilax paludinosus*). *Z. Säugetierk.*, 53: 358-364.

Baker, C. M. 1988. Vocalizations of captive water mongooses, *Atilax paludinosus*. *Z. Säugetierk.*, 53(2):83-91.

Baker, C. M. 1989. Feeding habits of the water mongoose (*Atilax paludinosus*). *Z. Säugetierk.*, 54:31-39.

Barré, N., Garris, G. I., Borel, G. & Camus, E. 1988. Hosts and population dynamics of *Amblyomma variegatum* (Acari: Ixodidae) on Guadeloupe, French West Indies. *J. Med. Entomol.*, 25(2):116-115.

Coblentz, B. E. & Van Vuren, D. 1988. Changes in kidney mass in the Indian mongoose (*Herpestes auropunctatus*). *J. Mammal.*, 69(3):614-618.

Desmet, K. & Hamdine, W. 1988. Densités de genettes (*Genetta genetta* Linné, 1758) en yeuseraie algérienne. *Mammalia* 52(4):604-607.

Enurah, L. U., Ocholi, R. A., Adeniyi, K. O. & Ekwonu, M. C. 1988. Rabies in a civet cat (*Civettictis civetta*) in the Jos Zoo, Nigeria. *Br. Vet. J.*, 144(5):515-516.

Enurah, L. U., Adeniyi, K. O., Ocholi, R. A., Spencer, T. H. I. & Badung, J. D. 1988. *Klebsiella pneumoniae* as a cause of pneumonia and septicemia in a civet kitten (*Civettictis civetta*) in the Jos Zoo, Nigeria. *J. Wildl. Dis.*, 24(3):585-586.

Grant, D. J. S. A. 1988. Confirmation of the occurrence of the marsh mongoose *Atilax paludinosus* in the Luangwa Valley, Zambia. *Black Lechwe* 9:31-32.

Hoppe-Dominik, Bernd. 1989. Premier recensement des grands mammifères dans le Parc National de la Marahoué en Côte-d'Ivoire. *Rev. Zool. Afr.*, 103(1):21-27.

Palomares, F. & Delibes, M. 1988. Time and space use by two common genets (*Genetta genetta*) in the Doñana National Park, Spain. *J. Mammal.*, 69(3):635-637.

Rood, J. P. 1988. What ever happened to the C Pack Sisters? *Nat. Hist.*, 97(2):40-47.

Taylor, M. E. 1988. Foot structure and phylogeny in the Viverridae (Carnivora). *J. Zool. (London)* 216(1):131-139.

Trivedi, K. K. & Gupta, S. P. 1987. Nematode parasites of vertebrates. 5. A new spirurid, *Spirocera skrjabini*, new species (Spiruridae Oerley, 1885) from a mongoose, *Herpestes vitticollis* Bennet, from Udaipur, Rajasthan, India. *Pak. J. Zool.*, 19(2):147-152.

Van Rompaey, H. 1988. *Osbornictis piscivora.* *Mammal. Species* 309:1-4.

Van Valkenburgh, B. 1988. Incidence of tooth breakage among large, predatory mammals. *Amer. Nat.*, 131(2):291-302.

# IUCN/SSC M&VSG membership

## Chairman

Mr Roland Wirth  
Franz-Senn-Strasse, 14  
8000 München  
FRG

## Members

Prof. R. Alagnac  
Parc Zoologique de Vincennes  
53, Avenue St. Maurice  
F-75012 Paris  
France

Mr Mahedi Andau  
Jabatan perhutanan  
Peti Surat. 311  
90007 Sandakan, Sabah  
Malaysia

Ms Merav Ben-David  
Dept. Zool. Tel Aviv University  
Ramat Aviv  
69978 Tel Aviv  
Israel

Dr B. Boeadi  
Curator, Dept. Mammals, Mus. Zool. Bogor  
Jl. Ir. H. Juanda, 3  
Bogor, Java Barat  
Indonesia

Mr A.-J. Braun  
Etude Loutre/Vison  
56370 Le Tour du Parc, Le Mare  
France

Dr G. B. Caddick  
Director, Breeding Projects Belize Zoo  
P. O. Box 474  
Belize City  
Belize

Dr Gerardo Ceballos  
Apartado Postal 23-D  
Toluca, Mex. 50120  
Mexico

Dr T. W. Clark  
Northern Rockies Conservation Coop.  
Box 2705  
Jackson, WY 83001  
USA

Dr M. Colyn  
104, Avenue de la Heronnière B 34  
1160 Bruxelles  
Belgium

Mr A. D. Cuarón  
Curator Mammals, Inst. Historia Natural  
Apto postal 6  
Tuxtla Gutierrez, Chiapas 29000  
Mexico

Dr Glyn Davies  
Dept. Anthropol., Univ. London  
Gowerstreet  
London WC1E 6BT  
UK

Shri R. K. Ghose  
Mammal & Osteo. Sect., Zool. Surv. India  
Lindsay Street, 8  
Calcutta 700087  
India

Dr C. P. Groves  
Dept. Prehist. & Anthropol. Austr. Ntl. Univ.  
G. P. O. Box 4  
Canberra ACT 2601  
Australia

Dr Robert Izor  
Curator, Div. Mammals, Field Mus. Nat.  
Hist.  
Roosevelt Road at Lake Shore Drive  
Chicago, IL 60605-2496  
USA

Dr Paul Joslin  
Int. Snow Leopard Trust  
16463 Southeast Thirty-fifth Street  
Bellevue, Washington 98008  
USA

Dr K. U. Karanth  
Centre for Wildlife Studies  
499, Kuvempu Nagar  
Mysore 570 023  
India

Dr G. U. Kurup  
Zool. Surv. India, West. Ghat Region St.  
2/355 Eranhipalam  
Calicut 673 006, Kerala  
India

Mr Tiit Maran  
Curator, Tallina Loomaaed  
Paldiski mnt., 145  
200035 Tallinn, Estland  
USSR

Prof. Rodney A. Mead  
Dept. Biol. Science, University Idaho  
Moscow, ID 83843  
USA

Mr Jarujin Nabhitabhata  
Curator, Thailand Inst. Sci. & Tech. Res.  
196 Phahonyothin Road, Bangkok  
Bangkok  
Thailand

Dr M. Nicoll  
WWF-Representation of Madagascar  
B. P. 4373  
101 Antananarivo  
Madagascar

Dr Gustav Peters  
Zool. Forsch. Inst. & Mus. A. Koenig  
Adenauerallee, 150-164  
D-5300 Bonn 1  
FRG

Dr R. A. Powell  
Dept. Zool., North Carolina State Univ.  
P. O. Box 7617  
Raleigh, NC 27695  
USA

Mr R. Ratajszczak  
Curator Wielkopolski Park Zoologiczny  
ul. Browarna, 25  
61-063 Poznan  
Poland

Mr Michael Riffel  
Hebelstrasse, 5  
D-7528 Karlsdorf-Neuthard 1  
FRG

Mr Paul Robinson  
Curator Southport Zoo  
Princes Park  
Southport, Lancashire  
UK

Mr Jerzy Romanowski  
Dept. Vert. Ecol., Inst. Ecology  
PL-05092 Dziekanow Lesny  
Poland

Dr D. A. Schlitter  
Curator Mammals, Carnegie Mus. Nat. Hist.  
5800 Baum Boulevard  
Pittsburgh, PA 15206  
USA

Mr Arnd Schreiber  
Zoologisches Institut, Heidelberg Univ.  
Im Neuenheimer Feld, 230  
D-6900 Heidelberg  
FRG

Prof. Rüdiger Schröpfer  
Universität Osnabrück, Biologie/Chemie  
Barbara Strasse, 11  
4500 Osnabrück  
FRG

Dr Mark E. Taylor  
Biol. & Environm. Consulting  
160 Woodcrest Avenue  
Toronto, Ontario M4J 3B9  
Canada

Dr E. T. Thorne  
Game & Fish Lab., University Wyoming  
P. O. Box 3312  
Laramie, WY 82071  
USA

Dr Harry Van Rompaey  
Jan Verberlei, 15  
2520 Edegem  
Belgium

Prof. Wang Yingxiang  
Head Dept. Vert. Taxon. Kunming Inst. Zool.  
Academia Sinica  
Kunming, Yunnan  
China

Dr Chris Wemmer  
Curator-in-charge, Natl. Zool. Park  
Conservation & Research Center  
Front Royal, VA 22630  
USA

Dr W. C. Wozencraft  
Div. Mammals, Natl. Mus. Nat. Hist.  
Smithsonian Institution  
Washington, DC 20560  
USA

## Subscriptions

Subscription for 1990 (two numbers planned) is 600 Belgian francs (approx. 15 US\$ or 10 £).  
Please make cheques payable to:

H. Van Rompaey  
Jan Verbertlei, 15  
2520 Edegem  
Belgium

or pay preferably by bank draft to:

Kredietbank  
Edegem Centrum  
2520 Edegem  
Belgium  
Account N° 419-0037021-92

When not paid in Belgium francs, costs for converting currency and banking may be high,  
please add 30% to cover these costs.

**This first number is sent free to all interested persons and institutions.  
The next numbers will only be sent to members of the  
IUCN/SSC Mustelid & Viverrid Specialist Group  
and to subscribers.**